

stryker



# Mako Partial Knee

Clinical summary



Volume 9

# Table of contents

---

<b>1. Introduction</b>	3
<b>2. What evidence is available for Mako Partial Knee</b>	4
2.1 Component placement accuracy	5
2.2 Surgical team learning curve	8
2.3 Soft tissue balance and bone preservation	9
2.4 Summary of evidence	10
<b>3. What are the potential clinical benefits of Mako Partial Knee</b>	11
3.1 Survivorship	11
3.2 Patient satisfaction	15
3.3 Clinical outcomes	16
3.4 Functional outcomes	19
3.5 Clinical outcomes lateral PKA	21
3.6 Continuum of care	22
3.7 Outcomes of patellofemoral arthroplasty	23
3.8 Outcomes of bicompartamental knee arthroplasty	24
3.9 Complication and revision	24
<b>4. Is Mako Partial Knee cost-effective?</b>	25
<b>5. Conclusion</b>	29
<b>6. References</b>	30

# 1. Introduction

Partial knee arthroplasty (PKA), also termed unicompartmental knee arthroplasty (UKA) when associated with a single compartment, has been performed for isolated single compartment knee osteoarthritis (OA) since the 1970s.<sup>1</sup> PKA can be carried out in the medial, lateral and/or patellofemoral (PF) compartments.

When compared to total knee arthroplasty (TKA), studies have shown that medial PKA patients experience greater retention of normal knee kinematics and accelerated recovery, while suffering less blood loss and reduced postoperative morbidity.<sup>2-5</sup> Lateral PKA is less common, comprising around one-eighth of all PKA cases.<sup>6</sup> However, lateral PKA has also been shown to be an effective treatment in the appropriate patient, with survivorship and outcomes comparable to medial PKA.<sup>6-8</sup> Patellofemoral arthroplasty has also demonstrated significant benefits to the patient when compared to TKA.<sup>9</sup> A 2018 double-blinded study showed that patients who underwent PF arthroplasty for isolated PF arthritis had a better overall knee-specific quality of life than patients who underwent TKA throughout the first two years after the operation.<sup>9</sup>

Despite the volume of evidence demonstrating the benefits of PKA, the procedure is known to be sensitive to surgical factors such as implant positioning and soft tissue balance.<sup>10</sup> This was highlighted in a study by Kazarian et al., where data from 253 medial PKA patients was retrospectively analyzed to assess the implant survival and radiographic outcomes after PKA, as well as the impact of component alignment and overhang on implant survival.<sup>11</sup> All procedures in the study were performed by two high-volume surgeons. The results showed that the incidence of PKA revisions and alignment outliers were greater than expected, even among high-volume arthroplasty surgeons performing an average of 14 PKAs per year.<sup>11</sup> Both alignment and overhang outliers were significant risk factors for implant failure.<sup>11</sup> The researchers emphasize that the ability of low-volume PKA surgeons to consistently attain accuracy in implant position is an important factor to investigate to help enhance PKA survivorship.<sup>11</sup>

The Mako System was introduced to assist surgeons in achieving accurate implant alignment, anatomic restoration and soft-tissue balancing, thereby supporting the surgeon in restoring native knee kinematics and enhancing patient outcomes.<sup>12-14</sup> This document summarizes the evidence to date that supports the use of Mako Robotic-Arm Assisted Surgery for PKA (Mako Partial Knee) including recent national joint registry data.<sup>15,16</sup>


## 2. What evidence is available on Mako Partial Knee?

Successful clinical outcomes following joint replacement are dependent on component placement and on restoring the natural kinematics of the knee. Component malalignment in PKA has been associated with stress concentrations, bone fracture and poor clinical outcomes.<sup>17-18</sup> The Mako System is designed to minimize the margin of error associated with component placement and to enhance the accuracy and reproducibility of PKA. Additionally, the Mako System helps enable the surgeon to dynamically balance soft tissue tensioning intraoperatively, with the goal of recreating natural knee kinematics. Clinical studies have shown that Mako Partial Knee has the potential to produce accurate and reproducible component placement in accordance with preoperative plans and to reestablish soft tissue balance.<sup>19-20</sup>

St Mart et al. (2022) conducted a prospective study to assess clinical and radiological outcomes for robotic- assisted unicompartmental knee arthroplasty (UKA, or PKA).<sup>21</sup> The study aimed to evaluate clinical and radiological outcomes following robotic-assisted PKA as well as any potential learning curves associated with the introduction of the technology.<sup>21</sup> The primary outcome measures were patient-reported outcome measures (PROMs) including Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC), Knee Society Score (KSS) and Oxford Knee Score (OKS), complications, implant survivorship, component positioning and learning curve.<sup>21</sup> Eighty-five patients comprising 100 knees were recruited and followed for  $21.0 \pm 4.3$  months.<sup>21</sup> At two years, there were significant and sustained improvements in PROMs.<sup>21</sup> A cumulative learning curve of 20 cases was noted.<sup>21</sup> The study concluded that robotic- assisted PKA achieved excellent implant accuracy to plan and clinical outcomes in the short-term.<sup>21</sup>

## 2.1 Component placement accuracy

A key clinical paper on Mako accuracy, published by Bell et al., reports on a randomized controlled trial (RCT) involving 120 patients.<sup>19</sup> The study compared patients who received robotic-arm assisted PKA (Restoris MCK n=62) with those who underwent manually implanted PKA (Oxford n=58).<sup>19</sup> Comparisons were made between groups in terms of the preoperative plan of femoral and tibial component positioning against the actual alignment achieved in three different planes (axial, coronal and sagittal).<sup>19</sup> Results showed more accurate component positioning in the robotic-arm assisted group, with lower root mean square (RMS) errors and significantly lower median errors in all six component parameters ( $p < 0.01$ ).<sup>19</sup> The proportion of patients with tibial slope within  $2^\circ$  of the target position was significantly greater using the robotic-arm assisted technique than the manual technique (80% compared with 22%,  $p = 0.0001$ ).<sup>19</sup> It was concluded that the Mako System more consistently placed the PKA implant in accordance with the preoperative plan (**Figure 1**).



**Figure 1.** Bell et al. (2016) showed that use of robotic-arm assisted PKA enabled surgeons to place the tibial and femoral components more accurately and consistently to plan.

**FS= Femoral Sagittal, FC=Femoral Coronal, FA= Femoral Axial, TS= Tibial Sagittal, TC\*= Tibial Coronal, TA=Tibial Axial.**

**\* = non-significant parameter.<sup>19</sup>**

These results were corroborated by a 2018 study performed by Kayani et al.<sup>22</sup> A single surgeon compared implant placement accuracy using radiographs from 60 consecutive conventional PKAs (Oxford) compared to the surgeon's first 60 consecutive Mako Partial Knees (Restoris MCK).<sup>22</sup> The Mako group had significantly ( $p < 0.001$ ) more accurate placement to plan for the femoral and tibial implants, as well as more accurate recreation of the knee's mechanical alignment, posterior tibial slope and joint line height.<sup>22</sup>

A study was performed at Washington University School of Medicine, U.S., by Kazarian et al. where postoperative radiological outcomes from 86 consecutive robotic-assisted PKAs (RA-PKA) were retrospectively reviewed and compared to 253 manual PKAs (M-PKA) drawn from a prior study at the same institute.<sup>23</sup> For the robotic-assisted group, 91.6% of all alignment measurements and 99.2% of all overhang measurements were within the target range.<sup>23</sup> All alignment and overhang targets were simultaneously met in 68.6% of RA-PKAs.<sup>23</sup> When comparing radiological outcomes between the RA-PKA and M-PKA groups, statistically significant differences were identified for rates of outliers in femoral coronal angle (2.3% vs. 12.6%;  $p = 0.006$ ), femoral sagittal angle (17.4% vs. 50.2%;  $p < 0.001$ ), tibial coronal angle (5.8% vs. 41.5%;  $p < 0.001$ ), and tibial sagittal angle (8.1% vs. 18.6%;  $p = 0.023$ ), as well as anterior (0.0% vs. 4.7%;  $p = 0.042$ ), posterior (1.2% vs. 13.4%;  $p = 0.001$ ), and medial (1.2% vs. 14.2%;  $p < 0.001$ ) overhang outliers.<sup>23</sup>

Matassi et al. considered the likelihood of robotic-assisted surgery in reducing the variability of coronal and sagittal component positioning between high- and low- volume surgeons.<sup>24</sup> A prospective cohort of 161 robotic-arm assisted medial PKA patients were divided into two groups: patients operated on by "high-volume" or "low-volume" surgeons.<sup>24</sup> They recorded intraoperative lower-limb alignment, component positioning, and surgical timing.<sup>24</sup> Postoperatively, they assessed coronal and sagittal femoral/tibial component alignment, ROM and patient-reported outcomes out to 1-year follow-up. Of the recruited knees, 149 ("high-volume": 101; "low-volume": 48) met inclusion.<sup>24</sup> No clinical difference in mechanical alignment nor coronal/sagittal component positioning were found ( $p > 0.05$ ) between groups. A significant difference was recorded in surgical timing ("high volume": 57 minutes; "low-volume": 86 minutes;  $p < 0.05$ ).<sup>24</sup> No superficial or deep infections or other major complications were developed during the follow-up. The study demonstrated that robotic-assisted surgery in medial PKA allowed for improved reproducibility with satisfactory clinical outcomes. Moreover, the study showed that differences in component positioning and lower limb alignment among low-and-high volume knee surgeons were almost eliminated.<sup>24</sup>

Gaggiotti et al compared the accuracy in implant positioning and limb alignment in a retrospective radiographic evaluation of Mako, which is a CT, image-based robot-assisted system used for RA-PKA (n=95) and an imageless robotic system (n=197).<sup>25</sup> Preoperative measurements were compared to postoperative conditions at one-year to evaluate hip-knee angle (HKA), posterior tibial slope (PTS), tibial component coronal alignment relative to Cartier's angle, and restoration of proper joint line (JL) height.<sup>25</sup> Both treatment groups exhibited improvement in HKA and PTS from baseline ( $p < 0.001$ ), but Mako's image-based RA-PKA group resulted in higher accuracy in implant positioning, as demonstrated by greater achievement of PTS within the target range of 2-8° (93.7% vs. 82.7% for image-based robotic vs imageless robotic, respectively;  $p = 0.01$ ), variations in Cartier's angle less than 3° (87.4% vs. 65% for image-based robotic vs imageless robotic, respectively;  $p < 0.001$ ) and changes in the height of the joint line less than 2mm (81.1% vs. 69.5% for image-based robotic vs imageless robotic, respectively;  $p = 0.04$ ).<sup>25</sup> Use of the Mako image-based robotic system was associated with a higher degree of accuracy in implant positioning and postoperative limb alignment as compared to an imageless robot-assisted system.<sup>25</sup>

Alongside placement accuracy, Kumar et al. published a study on the importance of correction of PKA postoperative angle.<sup>26</sup> The goal of the study was to understand the correlation between optimal postoperative varus deformity, correction angle and physiological constitutional varus deformity, and the influence of these parameters on WOMAC scores.<sup>26</sup> Robotic-assisted unicompartamental knee arthroplasty in 143 knees were retrospectively assessed with WOMAC scores preoperatively and after surgery at intervals up to 2 years.<sup>26</sup> The results showed that correction angles were more consistent in robotic than non-robotic outcomes and that preoperative varus and correction angles were well correlated.<sup>26</sup> Mean preoperative and postoperative varus deformities were 10.2° and 4.8°, respectively, with a mean correction angle of 5.4°. The preoperative varus and correction angles were found well correlated ( $r = 0.815$ ).<sup>26</sup>


However, the amount of improvement in the WOMAC total score was not correlated with postoperative varus angle.<sup>26</sup>

The results suggest that there was a significant improvement in the WOMAC score for patients at 1.5 months, 3 months, 6 months, 1 year and 2 years ( $p < 0.001$ ), however, the amount of improvement in the WOMAC total scores overall appeared not to be influenced by the degree of correction and that the correction angle is mostly related to the preoperative varus deformity.<sup>26</sup> More research was recommended.

## 2.2 Surgical team learning curve

In a Kayani et al study that included 60 consecutive conventional jig based UKAs and 60 consecutive robotic-arm assisted UKAs, RA-PKA demonstrated a learning curve of six cases for operating time and surgical team confidence levels equivalent to those of M-PKA.<sup>22</sup> The learning curve did not influence any of the associated accuracy variables, and accuracy to plan achieved with the Mako System was consistent between the surgeon's first Mako case and last 10 Mako cases.<sup>22</sup> This indicated that Mako Partial Knee surgery did not have a learning curve for accuracy in achieving the planned femoral and tibial implant position.<sup>22</sup> Further, no additional risk was observed for postoperative complications during the surgical team learning curve.<sup>22</sup>

Jinnah et al. have previously performed an extensive multicenter study to understand how learning curve may influence surgical time for Mako Partial Knee.<sup>27</sup> Eight hundred and ninety-two patients had a Mako Partial Knee performed by 13 different surgeons.<sup>27</sup> Surgical time was measured from insertion of the first bone pin to the acceptance of the final trial components. The average surgical time for all surgeons was  $56 \pm 20$  minutes.<sup>27</sup> The shortest average surgical time for an individual surgeon was  $38 \pm 9$  minutes and the longest was  $70 \pm 29$  minutes.<sup>27</sup> An average learning curve of 13 cases was proposed for the surgical time to reach a steady state (**Figure 2**).<sup>27</sup>



**Figure 2.** Typical Mako Partial Knee learning curve graph showing one surgeon's first 50 cases from a multicenter study by Jinnah et al. (2010). After approximately 13 cases, surgical time reached a steady state.<sup>27</sup>

Tay et al. concluded that the learning curve for Mako Partial Knee was 11 cases.<sup>28</sup> The study aim was to assess operating times, patient outcomes, limb alignment and component placement for both low and high-volume surgeons.<sup>28</sup> Five surgeons performed 152 consecutive robotic-arm assisted primary medial PKA, and measurements of interest were recorded.<sup>28</sup> Patient outcomes were measured with Oxford Knee Score, EuroQol-5D, and Forgotten Joint Score at six weeks, one year and two years.<sup>28</sup> The surgeons were divided into 'low' and 'high' usage groups based on total PKA (manual and robotic) performed per year.<sup>28</sup>

A learning curve of 11 cases was found, with a statistically significant decrease in operative time between the learning phase and the proficiency phase ( $p=0.01$ ), a significant improvement in femoral rotation between the two phases ( $p=0.02$ ), and more accurate insert sizing between the phases ( $p=0.03$ ), which highlighted areas that require care during the learning phase.<sup>28</sup> Despite decreased 6-week EQ-5D-5L Visual Analogue Scale (VAS) in the proficiency group (77 cf. 85,  $p < 0.01$ ), no difference was found in implant survival (98.2%) between learning and proficiency phases ( $p=0.15$ ), or between 'high' or 'low' usage surgeons ( $p=0.23$ ) at 36 months.<sup>28</sup> This suggests that the learning curve did not lead to early adverse effects in this patient cohort.<sup>28</sup> The conclusion was that Mako PKA showed learning curves for operative time (11 cases), and insert sizing, but not for implant survival at early follow-up.<sup>28</sup> The short learning curve regardless of PKA usage indicated that robotic-arm assisted PKA may be particularly useful for low-usage surgeons.

### 2.3 Soft tissue balance and bone preservation

From a soft tissue perspective, Plate et al. considered that the ability to effectively restore a patient's ligament length and tension may help with restoration of normal knee kinematics and muscle lever arms of the knee joint.<sup>20</sup> Their study examined the accuracy of dynamic, real-time ligament balancing for 52 Mako Partial Knees.<sup>20</sup> Gap distances at 0°, 30°, 60°, 90° and 110° flexion were assessed preoperatively and after final component implantation to establish whether ligament balancing was restored.<sup>20</sup> Ligament balancing was accurate up to 0.53 mm compared to the preoperative plan.<sup>20</sup> These results indicated the Mako System was capable of accurately and precisely reproducing the desired soft tissue balance.

In addition to this, a cadaveric investigation was carried out with the aim of quantifying the amount of bone preserved in medial RA-PKA compared to RA-TKA.<sup>29</sup> Eleven knees were selected and analyzed from seven cadavers. Results showed that RA-PKA procedures resected an average of  $11.6 \pm 1.33$  cm<sup>3</sup> (range: 9.85-13.7 cm<sup>3</sup>) whereas total knee procedures resected an average of  $59.7 \pm 9.65$  cm<sup>3</sup> (range: 47.4-78.3 cm<sup>3</sup>), demonstrating that for this study population, only 17% to 19% of the bone volume was resected in RA-PKA compared to robotic TKA.<sup>29</sup> The study highlighted that in robotic PKA, the femur preparation is contoured to match the implant, which may in turn contribute to enhanced bone preservation and retention of bone stock.<sup>29</sup>

In another study, Hampp et al. compared the extent of soft-tissue trauma sustained through Robotic-arm Assisted PKA (RA-PKA) and manual PKA (M-PKA).<sup>30</sup> Five surgeons, confident in RA-PKA and M-PKA techniques, were asked to prepare a total of 24 cadaveric knees.<sup>30</sup> Afterwards two independent surgeons were asked to estimate trauma to the patellar tendon, quadriceps tendon, anterior cruciate ligament (ACL), medial collateral ligament (MCL), medial capsule, posterior capsule, and posterior cruciate ligament (PCL) using a five-grade system where Grade 1 represented complete tissue preservation and Grade 5 represented over 76% tissue trauma.<sup>30</sup> When compared to the manual PKA group, RA-PKA had lower total trauma grading ( $p < 0.01$ ), lower posterior capsular damage ( $p < 0.01$ ), and less severe ACL damage ( $p < 0.01$ ).<sup>30</sup> The authors concluded that based on this analysis the use of Mako Partial Knee for PKA can result in significantly less soft tissue trauma compared to manual PKA.<sup>30</sup>

## 2.4 Summary of evidence

These studies demonstrated that Mako robotic-arm assisted technology equipped the surgeon to accurately and consistently place the femoral and tibial PKA components in accordance with preoperative plans, and to help the surgeon restore soft tissue balancing.<sup>19-20</sup> This technology is associated with a short learning curve to achieve time neutrality.

### 3. What are the potential clinical benefits of Mako Partial Knee?

Mako Partial Knee has been shown to deliver demonstrable clinical benefits.<sup>12-14</sup> Studies have investigated implant survivorship, patient satisfaction, clinical outcomes and functional outcomes in medial Mako Partial Knee, with favorable results in comparison to other surgical methods.<sup>12-14,31-32</sup> In lateral and PF Mako Partial Knee, encouraging clinical and functional outcomes have also been observed.<sup>9,33</sup> Furthermore, in both medial and lateral PKA, congruence of the nonsurgical and surgical compartments has been found to be restored, supporting the hypothesis that the resultant redistribution of contact forces across the patellofemoral joint could help address PF symptoms.<sup>34-35</sup>

#### 3.1 Survivorship


Favorable survivorship data was published by Roche et al.<sup>37</sup> This site performed a retrospective review of prospectively collected data in their institution's registry on patients who underwent medial RA-UKA.<sup>37</sup> The final query consisted of 185 patients. Patients had a mean age of 65 years and mean body mass index (BMI) of 31.6 kg/m<sup>2</sup> with a mean follow-up of 10 years.<sup>37</sup> Ten-year survivorship of the study cohort demonstrated 99% survivorship at 10 years following the index procedure.<sup>37</sup> Majority of the patients (97%) stated they were either "very satisfied" (80.95%) or "satisfied" (16.19%) with the outcomes of their procedure.<sup>37</sup> Two patients were revised during the study period. This study represents one of the few long-term analyses of patients undergoing medial RA-UKA to report on long-term survivorship and patient-satisfaction.<sup>37</sup>

In a U.S. study published by Burger et al. midterm implant survivorship was evaluated for RA- PKA patients.<sup>38</sup> The research involved a retrospective review of patients who underwent robotic-arm assisted PKA between 2007 and 2016.<sup>38</sup> Study participants received a fixed-bearing medial or lateral PKA, patellofemoral arthroplasty (PFA), or bicompartamental knee arthroplasty (involving a PFA plus medial PKA), and the mean follow-up was 4.7 years (2.0 to 10.8).<sup>38</sup> The five-year survivorship rate of medial PKA (n=802), lateral PKA (n=171) and PFA/bicompartamental knee arthroplasty (n=35/10) was 97.8%, 97.7% and 93.3%, respectively.<sup>38</sup>

Comparable data was previously confirmed in a multicenter longitudinal study evaluating short- and midterm survivorship of robotic-arm assisted medial PKA, which demonstrated 98.8% survivorship (in 909 knees) at 2.5-year follow-up and 97% survivorship (in 432 knees) at 5.5-year follow-up.<sup>12,39</sup> This survivorship rate was greater than rates derived from high-volume surgeon data and registry data for conventional PKA (**Figure 3**).<sup>12,39</sup>

These studies concluded that the favorable survivorship observed resulted from Mako's ability to help enable surgeons to achieve more accurate component positioning when compared to implant placement using manual techniques.<sup>12,39</sup>

When considering patient satisfaction, 91% of patients without revision reported being either very satisfied or satisfied at 10.2-years.<sup>39</sup>



**Figure 3.** Survivorship data from Pearle et al. (2017)<sup>39</sup> and Kleebblad et al. (2018)<sup>12</sup> on robotic-arm assisted PKA compared to studies in literature and annual registries reporting 2 to 3 years and 5 to 6 years conventional PKA survivorship data.

An RCT by Gilmour et al. comparing patients who underwent medial Mako Partial Knee (Restoris MCK) with those who underwent manual medial PKA (Oxford) demonstrated favorable results for Mako PKAs.<sup>40</sup>

Specifically, Mako Partial Knee patients had 100% survivorship compared to 96.3% in the manual group at two years post-operation.<sup>40</sup> The 100% survivorship rate was maintained in the Mako Partial Knee group at Five years and Ten years post-operation.<sup>41,42</sup>

Bayoumi et al. published 10-year survivorship and satisfaction results in 2023 based on data from a multicenter study.<sup>43</sup> The multicenter longitudinal study evaluated long-term survivorship of robotic-arm assisted medial UKA patients who underwent the index procedure between March 2009 and December 2011.<sup>43</sup> A total of 366 patients (411 knees) were analyzed with a mean follow-up of  $10.2 \pm 0.4$  years, which demonstrated 98.8% survivorship at 2.5-year follow-up (n = 909), 97% survivorship at 5.5-year follow-up (n = 432), and 92.2% survivorship at 10.2-year follow-up.<sup>43</sup>

These findings were also reflected in a study by Zambianchi et al, who examined the incidence of revision on 188 Mako Partial Knee cases in 196 patients from a single center at a mean follow-up of 11.1 years (range 10.0-11.9 years).<sup>44</sup> Seven patients underwent revision procedures, resulting in a survivorship rate of 96.4% (95% confidence intervals 94.6-99.2).<sup>44</sup> This cohort also has Forgotten Joint Scores (FJS) assessed. The mean FJS was 82.2 (standard deviation 23.9).<sup>44</sup> Additionally, 88.8% of patients achieved the Patient Acceptable Symptoms State (PASS) of FJS greater than 40.63.<sup>44</sup>

Registry data recorded in the Australian Orthopaedic Association National Joint Replacement Registry (AOANJRR) is available for the Restoris MCK partial knee system, which is implanted using Mako Partial Knee. Data included in an industry report (i.e. available to industry on-demand) generated in June 2024 shows a revision rate of 6.0% (CI: 5.2 – 6.9) at nine years.<sup>52</sup> This report also presents data as revisions per 100 observer years (Revs/100 Obs. Yrs.), with the Restoris MCK showing a rate of 0.82 Revs / 100 Obs. Yrs. (CI: 0.74 – 0.91). These data do not distinguish between implantation for medial or lateral implantation as the same components are used.<sup>52</sup>

Similarly, the Restoris MCK partial knee system appears in the National Joint Replacement Registry of England and Wales (NJR)<sup>16</sup> report in 2024. Restoris MCK currently has an ODEP rating of 7A\* with a revision rate of 2.24%.<sup>16</sup>


### **Midterm survivorship for isolated lateral compartmental arthritis**

Midterm survivorship for robotic-assisted PKA was investigated by Heckmann et al.<sup>45</sup> This retrospective case series assessed results of all lateral PKAs performed by a single surgeon between 2013 and 2019.<sup>45</sup> Patient demographics, surgical variables and Kozinn and Scott criteria were collected, and implant survivorship was estimated using the Kaplan-Meier method with the endpoint being conversion to TKA.<sup>45</sup>

The results identified 84 lateral PKAs which met the inclusion criteria with a mean follow-up of 4.0 years (range 2.0-7.0).<sup>45</sup> Five-year survivorship was 92.9% (95% confidence interval [CI] 84.5-96.7) with all cause revision and 100% conversion to TKA as an endpoint.<sup>45</sup> Six-year survivorship was 88.9% with one additional patient being revised to TKA after five years. The average Forgotten Joint Score was 82.7% and patient satisfaction 4.7/5.<sup>45</sup>

The study reported high midterm survivorship and excellent patient-reported outcomes and concluded that robotic-assisted lateral PKA is a viable treatment option for isolated lateral compartment arthritis.<sup>45</sup>

The revision rates for Mako Partial Knee with Restoris MCK have been published in cohort studies, economic analyses, level I clinical trials (RCTs) and international registries (**Figure 4**).<sup>12, 38, 39, 42-44</sup> The evidence supports excellent survivorship of the Restoris MCK implant used in both the medial and lateral compartment (**Table 1**).<sup>2, 38, 39, 42-44</sup>



**Figure 4.** Graph indicating Mako Partial Knee revision rates with data taken from cohort studies, economic analyses, level I clinical trials (RCTs) and international registries.

Study	Months	n	CRR
Cool et al. <sup>46</sup>	24	246	0.81
Pearle et al. <sup>39</sup>	29.6	909	1.2
St Mart et al. <sup>47</sup>	36	2851	2.6
Dretakis et al. <sup>48</sup>	36	51	0
Deese et al. <sup>49</sup>	54	81	1.2
Banger et al. <sup>41</sup>	60	65	0
Burger et al. <sup>38</sup>	60	802	2.2
NJR England and Wales <sup>16</sup>	60	2919	2.24
Kleebblad et al. <sup>12</sup>	68.4	384	3
Zambianchi et al. <sup>44</sup>	70.8	216	2.2
AOANJRR Industry report <sup>152</sup>	108	11076	6.0
Roche et al. <sup>37</sup>	120	185	2
Blyth et al. <sup>42</sup>	120	65	0
Bayoumi et al. <sup>43</sup>	122.4	411	7.9
Zambianchi et al. <sup>44</sup>	133.2	196	3.6
AOANJRR All Uni (2024) <sup>15</sup> (for reference)	12	-	1.9
AOANJRR All Uni (2024) <sup>15</sup> (for reference)	36	-	4.4
AOANJRR All Uni (2024) <sup>15</sup> (for reference)	60	-	6.1
AOANJRR All Uni (2024) <sup>15</sup> (for reference)	120	-	11.3


**Table 1.** Survivorship

## 3.2 Patient satisfaction

In two longitudinal clinical trials, most Mako Partial Knee patients were “very satisfied” or “satisfied” with their joint replacement.<sup>12,39</sup> These studies performed follow-up at 2.5 years (909 knees) and 5.5 years (432 knees) respectively with patients who underwent medial Mako Partial Knee procedures.<sup>12,39</sup> 92% of patients reported satisfaction with their knee 2.5 years postoperatively and 91% of patients reported satisfaction at 5.5 years (**Figure 5**).<sup>12,39</sup> For context the Swedish Knee Arthroplasty Registry reports 83% of 7,860 patients who underwent manual medial PKA were satisfied with their knee at an average six-year follow-up.<sup>50</sup>

Using the Mako System, Coon et al. performed 152 (71.3%) medial RA-PKAs, 33 (15.5%) lateral RA-PKAs, 20 (9.4%) medial bicompartamental RA-PKAs and 8 (3.8%) patellofemoral RA-PKAs.<sup>51</sup> All surgical procedures had high patient satisfaction with an average of 82.5% of patients reporting being very satisfied or satisfied at six months, which increased to 89.5% at two years.<sup>51</sup> The lateral RA-PKA group reported 100% satisfaction two years post-operation.<sup>51</sup>


Comparable midterm patient satisfaction data was published by Burger et al in a large single-surgeon study of 1018 knees, where a large proportion of patients who underwent robotic-assisted PKA reported high satisfaction levels.<sup>38</sup> The mean follow-up was 4.7 years (2.0 to 8.0), and the results showed that 90.7% of medial RA-PKA patients, 92.6% of lateral RA-PKA patients and 78.9% of RA-PFA or bicompartamental knee arthroplasty patients were either very satisfied or satisfied with their knee function.<sup>38</sup>



**Figure 5.** Midterm patient satisfaction with medial Mako Partial Knee procedures (Kleebblad et al., 2018<sup>12</sup> and Pearle et al., 2017<sup>39</sup>).

### 3.3 Clinical outcomes


Analysis by Blyth et al<sup>13</sup> of data from a prior RCT (Bell al.<sup>19</sup>) found that patients who underwent medial Mako Partial Knee experienced less pain than those who underwent manual surgery during the 90-day postoperative period.<sup>13</sup> Median pain scores were 55.4% lower in robotic-arm assisted patients compared to manual patients from day one to day 56 (**Figure 6**).<sup>13</sup> Furthermore, the robotic-arm assisted patients had a better American Knee Society Score (AKSS) at three months postoperatively and at one year postoperatively, and a greater proportion of robotic-arm assisted patients showed improvements in their UCLA Activity Score.<sup>13</sup> Through binary logistic regression, the study was also able to predict the key factors associated with achieving excellent outcomes on the AKSS. These factors were a preoperative UCLA Activity Score level >5 and the use of robotic-arm assisted surgery, although these do not withstand adjustment for multiple comparisons.<sup>13</sup>



**Figure 6.** Visual analog pain score collected up to 90 days postoperatively in a RCT of manual vs. robotic arm-assisted medial PKA procedures.<sup>13</sup>

In two separate studies, evidence showed that medial Mako Partial Knee patients were more likely to “forget” their artificial joint during daily life compared to those who underwent manual TKA.<sup>31-32</sup> Zuiderbaan et al. administered The Forgotten Joint Score (FJS) questionnaire one and two years postoperatively.<sup>31</sup> Scores were compared between 65 patients who underwent medial Mako Partial Knee and 65 patients who underwent manually instrumented TKA.<sup>31</sup> Results demonstrated patients who underwent medial robotic- arm assisted PKA were more likely to forget their artificial joint in daily life. Similarly, in a separate powered cohort study from the U.K., conducted by Clement et al.<sup>32</sup> Thirty patients who underwent Mako Partial Knee were propensity score matched to 90 patients who underwent manual TKA for isolated medial compartment arthritis.<sup>32</sup> The findings from this study showed that the six-month FJS was significantly greater for the RA-PKA group compared to the manual TKA group (difference 24.2,  $p < 0.001$ ) (**Figure 7**).<sup>32</sup>

### Forgotten joint score



**Figure 7.** FJS at six months, one and two years post-operation showing significantly higher scores in the medial Mako Partial Knee group ( $p < 0.001$ ,  $p = 0.002$  and  $p = 0.004$ , respectively).<sup>31-32</sup>

Furthermore, the same powered (1:3 ratio) cohort study by Clement et al. published early postoperative outcomes data, where statistically and clinically significant greater knee-specific functional outcomes were observed in RA-PKA patients compared to those who underwent manual TKA.<sup>32</sup> Findings showed that the RA-PKA group had a significantly greater six-month Oxford Knee Score by nearly eight points, and there was a five-point (95% confidence interval 1.9 to 8.1;  $p < 0.001$ ) greater improvement from the respective pre-operative baseline in the RA-PKA group compared to the manual TKA group, which was greater than the minimal clinically important difference.<sup>32</sup> This positive early outcome data was further fortified within the study as the researchers also found that the RA-PKA group had significantly better postoperative pain visual analogue scale (VAS) scores compared with the manual TKA group (**Table 2**).<sup>32</sup>

Mean PROM (SD)	rUKA	mTKA	Difference (95% CI)	p-value*
Postoperative OKS	44.2 (4.4)	36.5 (9.4)	7.7 (4.2 to 11.3)	<0.001
Postoperative FJS	77.1 (25.9)	52.9 (32.6)	24.2 (11.2 to 37.2)	<0.001
Postoperative EQ-5D	0.913 (0.126)	0.764 (0.248)	0.148 (0.054 to 0.241)	0.002
Postoperative pain VAS	93.6 (12.3)	76.4 (24.8)	20.5 (9.9 to 31.0)	<0.001

\*Unpaired t-test.

CI, confidence interval; EQ-5D, EuroQol five-dimension questionnaire; FJS, Forgotten Joint Score; M-TKA, manual total knee arthroplasty; OKS, Oxford Knee Score; PROM, patient-reported outcome measure; rUKA, robotic unicompartmental knee arthroplasty; VAS, visual analog scale.

**Table 2.** Six-month postoperative outcome measures and differences between robotic PKA vs. manual TKA.<sup>32</sup>


Zambianchi et al. published long-term data assessing clinical outcomes following medial robotic-arm assisted PKA at a minimum of ten years.<sup>44</sup> A total of 188 patients (196 knees) were evaluated at a mean follow-up of 11.1 years (range 10.0 – 11.9 years), resulting in a mean FJS-12 of 82.2, which aligns well with the shorter-term data published by Zuiderbaan and Clement.<sup>31,32,44</sup> The ten-year FJS-12 outperformed the five-year results from the same study cohort, published previously,<sup>52</sup> by an average of 7 points, which the authors hypothesized to be attributed to lower physical demands that accompany aging that may in turn lead to reduced joint awareness.<sup>44</sup>

Overall, results of these studies suggested positive clinical and patient-reported outcomes of robotic-arm assisted medial, lateral PF and bicompartmental PKA.

### 3.4 Functional outcomes

Gait analysis has been used to compare outcomes of robotic-arm assisted PKA patients to those of manual Oxford PKA patients. In an RCT, Motesharei et al. compared the gait of 31 RA-PKA patients to 39 Oxford PKA patients one year postoperatively.<sup>53</sup> Both groups were compared to a control group of 50 healthy subjects obtained from the University of Strathclyde's archive.<sup>53</sup>

Results from this study showed statistically significant differences in knee joint kinematics during level walking between the robotic-arm assisted and manual PKA groups.<sup>53</sup> The robotic-arm assisted group achieved a higher knee excursion (18.0°, SD 4.9°) compared to the manual group (15.7°, SD 4.1°).<sup>53</sup> There was no significant difference between the healthy group and the robotic-arm assisted group, but there was a significant difference between the healthy group and the manual group ( $p < 0.001$ ) (**Figure 8**).<sup>53</sup>



**Figure 8.** Mean knee excursion angles of the control group, the robotic-arm assisted and manual PKA groups during the stance phase of gait at one year post-operation.<sup>54</sup>

This study was repeated at five-years postoperatively by Millar et al., though on fewer patients (25 Mako vs. 21 Oxford), and the differences seen at one year were maintained.<sup>14</sup> Results showed that the Mako group achieved significantly greater knee flexion in weight acceptance than the conventional group.

These findings suggested that the potential for improved alignment offered by the Mako System may result in enhanced function of the knee during gait. These findings also demonstrated that the use of the Mako System resulted in a gait pattern that facilitated the normal function of the knee more closely than the conventional technique.<sup>14,53</sup>

A clinical study by Borus et al. assessed functional performance in patients who received robotic-arm- assisted PKA compared to those who received manual TKA.<sup>54</sup> Tests included a six-minute walk, timed up and go, and stair ascend/descend, which were measured preoperatively at six weeks and at three months postoperatively.<sup>54</sup> Although a statistically significant difference in functional performance change between groups was not reached, the authors highlighted that at six weeks, the RA-PKA group was able to walk an additional 21.00 meters (68.90 feet) compared to just 5.95 meters (19.52 feet) for the manual TKA group.<sup>54</sup> Remarkably similar functional differences were observed with the timed up and go and stair ascend/descend tests, suggesting that robotic PKA provided functional benefits that were at least equivalent to manual TKA.<sup>54</sup>

Research by Coon et al. on medial, lateral, medial bicompartamental and patellofemoral RA-PKAs carried out with Mako Partial Knee showed that at two years postoperatively, 87.9% of patients were as active or the same as they expected they would be before surgery.<sup>55</sup>

In addition, the average distance walked at discharge was 79.8 meters, and 90.9% of patients were walking without support three weeks postoperatively.<sup>55</sup> Lastly, 65 patients were employed at time of surgery, and 86% of those patients returned to work six weeks after their operation.<sup>55</sup>

### 3.5 Clinical outcomes lateral PKA

Lateral PKA is less frequently performed within the general population, accounting for just one-eighth of PKA cases.<sup>6</sup> However, this procedure has been shown to be effective for the appropriate patient, achieving reliable improvements in pain, function and implant survivorship.<sup>6-8</sup> The Mako robotic platform offers potential benefits through its demonstrated accuracy and reproducible implant positioning, helping to minimize the margin of error associated with component placement.<sup>45</sup> In addition, the platform enables intraoperative dynamic soft tissue balancing to help the surgeon recreate the patient's natural knee kinematics.

Positive results have been reported by several studies examining lateral Mako Partial Knee. A retrospective study conducted by van der List et al. compared two-year postoperative functional outcomes using the Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC score) and FJS, between 143 medial and 36 lateral Mako Partial Knee procedures.<sup>33</sup> Equivalent functional outcomes were noted for both medial and lateral PKA procedures.<sup>33</sup>

Similar positive survivorship data was published by Augart et al.<sup>56</sup> The authors performed a search of their institution's joint registry and found 88 lateral robotic- arm assisted PKA patients, with a mean follow-up of 24.4 months  $\pm$  10.7 months, who had 100% survivorship at final follow-up without revision to TKA.<sup>56</sup> The positive data observed thus far from medial and lateral Mako Partial Knees suggests that the potential benefits offered by the Mako robotic platform, with regards to surgical planning, precision, reproducibility and intraoperative soft tissue adjustments, have the potential to help enhance surgical accuracy during these technically demanding procedures.<sup>56</sup>

### 3.6 Continuum of care

As mean patient age decreases, partial knee arthroplasty is often considered as a conservative treatment to delay need for a total knee replacement. Studies of joint line restoration, patella tracking, and medial and lateral compartment congruency have been conducted at Hospital for Special Surgery in New York.<sup>34-36</sup> In all three studies, congruence of the surgical compartment was restored through the Mako procedure and implant. Congruence and joint line of the nonoperative compartment were also restored ( $p=0.001$ ).<sup>34</sup> The authors hypothesized that the improved patellofemoral congruence after Mako Partial Knee may lead to redistribution of contact forces across the patellofemoral joint and secondarily treat PF symptoms.<sup>34</sup>

Similarly, in a 2020 study by Burger et al., researchers aimed to explore the effect of patellofemoral joint pathology on lateral RA-PKA.<sup>57</sup> In particular, the effect of preoperative radiological degenerative changes and alignment on patient-reported outcome measures (PROMs) after lateral RA-PKA was evaluated, as well as the influence of lateral RA-PKA on the alignment of the patellofemoral joint.<sup>57</sup> A consecutive series of 140 knees in 130 patients who underwent Mako robotic-arm assisted fixed-bearing lateral RA-PKA were retrospectively reviewed.<sup>57</sup> Radiological evaluation was conducted to obtain a Kellgren Lawrence (KL) grade, an Altman score and alignment measurements for each knee. Postoperative PROMs were assessed using the Kujala (Anterior Knee Pain Scale) score, Knee Injury and Osteoarthritis Outcome Score Joint Replacement (KOOS JR) and satisfaction levels.<sup>57</sup> The results showed that at mean 4.1 years (2.0 to 8.5) follow-up, good to excellent Kujala scores were reported, and the presence of mild to moderate preoperative patellofemoral joint osteoarthritis had no impact on these scores (KL grade 0 vs. 1 to 3,  $p = 0.203$ ; grade 0 to 1 vs. 2 to 3,  $p = 0.674$ ).<sup>57</sup> Comparable scores were reported by patients with osteoarthritis evident on either the medial or lateral patellofemoral joint facet, and patients with abnormal patellar congruence and tilt angles ( $\geq 17^\circ$  and  $\geq 14^\circ$ , respectively) reported good to excellent Kujala scores.<sup>57</sup> Furthermore, it was evident that lateral PKA resulted in improvements to patellofemoral alignment.<sup>57</sup> The findings from this study demonstrate that mild to moderate preoperative radiological degenerative changes and malalignment of the patellofemoral joint may not be associated with poor patient-reported outcomes at midterm follow-up after lateral fixed-bearing PKA.<sup>57</sup>

The researchers went on to suggest that this may be explained by realignment of the patella and the resulting redistribution of loads across the patellofemoral joint.<sup>57</sup>

### 3.7 Outcomes of patellofemoral arthroplasty

The purpose of patellofemoral arthroplasty (PFA) is to address the pain caused at the patellofemoral joint without performing a more substantial total knee surgery that would sacrifice additional bone. However, past literature has reported conflicting success rates of PFA as a surgical treatment for patellofemoral OA.<sup>58</sup> Odgaard et al. used a multicenter, double-blinded RCT to compare clinical outcomes associated with PFA and TKA to establish whether there was an advantage to either option.<sup>9</sup> They found that PFA patients recovered quicker than TKA patients, and the functional outcomes were also better for PFA patients.<sup>9</sup> The average TKA patient lost almost three months of knee function postoperatively during the first two years, relative to the PFA patient.<sup>9</sup> It was concluded that PFA was a superior option to TKA in patients with patellofemoral OA.<sup>9</sup>

Encouraging functional data was observed in another study by Noyes et al. which examined 44 consecutive patients in a single-surgeon series who underwent RA-PFA (51 knees, seven bilateral).<sup>59</sup> The authors analyzed both sports and work activity levels in younger active patients with a mean age of 37.2 (range, 21-50 years) enrolled over the course of ten years (2009-2018) with a mean follow-up of 5.3 years (range 2-9.3 years).<sup>59</sup> All patients received a comprehensive clinical evaluation, Cincinnati Knee Rating System (CKRS) and patient psychometric ratings of the substantial clinical benefit (SCB) and the Patient Acceptable Symptom State (PASS).<sup>59</sup> They also received radiographic evaluation. Results showed high levels of participation in low-impact sports, rising from a baseline of 20% participation preoperatively, increasing to 80% postoperatively with an additional 7% able to perform higher-impact jumping-pivoting sports ( $p < 0.001$ ).<sup>59</sup> Pre-operatively, patients reported moderate to severe knee pain with activities of daily living in 91% of knees, which reduced to 11% at follow-up.<sup>59</sup> SCB rating demonstrated that 87% of patients rated their knee as good, very good, or normal.<sup>59</sup> PASS analysis showed 89% of patients were satisfied with the overall surgical outcome and 93% would be willing to undergo surgery again. In total, five knees underwent conversion to TKA (10% failure), including one knee that experienced a post-operative joint infection that required implant removal and subsequent total knee replacement.<sup>59</sup> This research demonstrated that robotic-arm assisted PFA was a successful treatment option in younger active patients with isolated PF arthritis, enabling a high return to low-impact recreational activities and occupations.<sup>59</sup>

### 3.8 Outcomes of bicompartamental knee arthroplasty

Bicompartamental knee arthroplasty (BiKA) may be an alternative for TKA candidates with localized arthritis. One advantage of BiKA in comparison to TKA is that it is less invasive.<sup>60</sup> It requires less bone removal and preserves the anterior and posterior ligaments which may lead to better stability and proprioception for the patient.<sup>60</sup> Studies have reported BiKA resulting in less blood loss, shorter hospital stay, and faster rehabilitation when compared to TKA.<sup>61</sup> However, there are also concerns around the complication rates for BiKA.

Gaudiani et al. published on their prospectively maintained cohort of 50 patients (53 knees) who underwent Robotic-Arm Assisted BiKA (RA-BiKA: patellofemoral and medial compartment) at five- and seven-year follow-up.<sup>62</sup> The group reported high survivorship rates, with 96% at five years and 93% at seven years.<sup>62</sup> At a mean follow-up of 7.1 years (range 7.0 – 7.3), 89% of patients reported being either satisfied or neutral with their RA-BiKA where 11% reported being not satisfied.<sup>62</sup> A mean Knee Society – Function Score of  $80.5 \pm 15.8$  with 82% of patients reporting walking more than 10 blocks, 89% walking without support, and 100% able to go up and down stairs with 61% requiring use of a handrail.<sup>62</sup>

### 3.9 Complication and revision

In 2022 Sun et al. conducted a systematic review and meta-analysis to compare complication and revision rates of RA-PKA vs. M-PKAs.<sup>63</sup> They considered eligible case-control studies comparing robotic-assisted and manual PKA with data from all eligible articles independently extracted by two authors.<sup>64</sup> Differences in outcomes between M-PKA and RA-PKA were analyzed by calculating the corresponding 95% CIs and pooled relative risks (RRs).<sup>63</sup>

The results included in the final meta-analysis were taken from a total of 16 studies involving 50,024 patients.<sup>63</sup> It was found that RA-PKA had fewer complications (RR: 0.52, 95% CI: 0.28 to 0.96,  $p=0.036$ ) and lower revision rates (RR: 0.42, 95% CI: 0.20 to 0.86,  $p=0.017$ ) than M-PKA.<sup>63</sup> No significant differences in non-implant-specific complications between the two surgical techniques (RR: 0.80, 95% CI: 0.61 to 1.04,  $p=0.96$ ) were observed. No publication bias was found in this meta-analysis.<sup>63</sup>

## 4. Is Mako Partial Knee cost-effective?

With rising demand for PKA in patients who seek restored function and a quicker recovery time, a U.S. study performed by Kazarian et al. evaluated the cost-effectiveness of RA-PKA compared to TKA as well as nonsurgical treatment (NST).<sup>64</sup> Using a Markov decision analytic model, the authors assessed lifetime costs and QALYs as function of Age at Time of Initial Treatment (ATIT) of patients with end-stage unicompartmental knee osteoarthritis.<sup>64</sup> The analysis included direct medical and indirect costs. Models for ATITs at five-year intervals from age 40 through 90, and RA-PKA was found to be more cost-effective compared to NST for patients aged 40 to 86.<sup>64</sup> Furthermore, when surgical treatments were compared, RA-PKA dominated TKA by generating more QALYs than TKA for all ATITs.<sup>64</sup> The authors further concluded that if PKAs were performed as 12% to 20% of the total volume of knee arthroplasties versus the less than 8% observed, it would lead to a lifetime cost-savings of 987 million to 1.5 billion U.S. dollars and increased lifetime QALY accumulation of 124,403 to 217,705 across the U.S. population.<sup>64</sup>

In a separate U.K.-based study, a Markov decision analysis by Clement et al. was performed to assess the cost-effectiveness of RA-PKA relative to manual TKA (M-TKA) and manual PKA (M-PKA) for patients with isolated medial compartment OA of the knee with a mean age of 65 years.<sup>65</sup> The study objective was to identify the cost per quality adjusted life-year of RA-PKA relative to M-TKA and M-PKA. Model inputs included hospital costs, implant survival and mortality rate. Using a model with an annual case volume of 100 patients, the cost per QALY of M-PKA was £1,395 and £1,170 relative to M-TKA and M-PKA, respectively.<sup>65</sup> The cost per QALY was influenced by case volume: a low-volume center performing 10 cases per year would achieve a cost per QALY of £7,171 and £8,604 relative to M-TKA and M-PKA, respectively.<sup>65</sup> For a high-volume center performing 200 rPKAs per year with a mean two-day length of stay, the cost per QALY would be £648; if performed as day cases, the cost would be reduced to £364.<sup>65</sup> For a high-volume center performing 200 RA-PKAs per year with a shorter length of stay of one day relative to M-PKA, the cost per QALY would be £574.<sup>65</sup> Furthermore, the cost per QALY of RA-PKA decreased with reducing length of hospital stay and with increasing case volume, compared with M-TKA and mPKA.<sup>65</sup> The model showed that RA-PKA was a cost-effective alternative to M-TKA and M-PKA for patients with isolated medial compartment OA of the knee.

Varughese et al retrospectively compared the first 120 RA-PKA patients to 120 patients who underwent M-TKA by four high-volume surgeons at a single center.<sup>66</sup> RA-PKA resulted in a significantly shorter length of stay compared to M-TKA (1.8 vs 4.8 days;  $p < 0.001$ ) and RA-PKA patients were discharged home more frequently compared to M-TKA (98.3% vs 74.2%;  $p < 0.001$ ).<sup>66</sup> Operative time was shorter for M-TKA vs RA-PKA, (75.9 min vs 86.0 min;  $p = 0.004$ ). Surgery-related readmission within 90-days was similar between the groups (0.8% RA-PKA vs 1.7% M-TKA,  $p = 1.0$ ) and both occurred at such a low rate that these costs were not included in the economic analysis.<sup>66</sup> RA-PKA resulted in a lower average cost per patient than M-TKA (AUD\$12,020 vs AUD\$19,199), shorter length of hospital stay, and reduced implant, consumable, sterilization and in-hospital rehabilitation costs.<sup>66</sup> Additionally, there was statistically lower use of opioids between RA-PKA and M-TKA (125.0 Morphine Equivalents (ME) vs 522.1 ME,  $p < 0.001$ ) during the 90-day post-operative period and reduced use of opioids per day for RA-PKA versus M-TKA (66.3 ME/day vs 114.7 ME/day,  $p < 0.001$ ).<sup>66</sup>

In summary, these models demonstrated that in patients with isolated medial compartment arthritis, RA-PKA was observed to be a more cost-effective procedure compared to nonsurgical treatment and M-TKA for the specified age groups modelled, thus concluding RA-PKA was cost effective compared to M-TKA.

In the U.S., in a study by Cool et al., reasons for revisions and associated costs were analyzed for unicompartmental arthroplasty cases.<sup>46</sup> UKA procedures were identified using a commercial administrative claims database to evaluate hospital admissions for revision surgeries.<sup>46</sup> Robotic UKA (rUKA, Mako Partial Knee) and manual UKA (mUKA, manual partial knee) procedures performed between March 1, 2013 and July 31, 2015 were used to calculate the rate of revisions within 24 months of the index procedure.<sup>46</sup>

Cases were propensity matched 2:1 based on age, sex, race, geographic division, high-cost comorbidities and concentration of healthcare specialists per 100,000 population to control for outside confounding factors at case index.<sup>46</sup> A total of 738 commercial health plan patients (246 RA-UKA, 492 M-UKA) were selected for inclusion in the analysis.<sup>46</sup> Results indicated fewer revision procedures in RA-UKA (0.81% (2/246) vs. 5.28% (26/492);  $p = 0.0017$ ) and RA-UKA patients incurred lower mean costs for the index stay plus revision(s) (\$26,001 vs. \$27,977;  $p > 0.05$ ).<sup>46</sup> Lower length of stay at index was also noted in the RA-UKA group (1.77 vs. 2.02 days;  $p = 0.0047$ ).<sup>46</sup> The study concluded that patients who underwent RA-UKA had fewer revision procedures, shorter LOS and incurred lower mean costs at 24 months.<sup>46</sup>

Findings from a 2020 U.K. cohort study involving 30 Mako Partial Knees compared to 90 propensity-matched Manual TKAs (M-TKA) showed that the length of stay was significantly ( $p < 0.001$ ) shorter in the RA-PKA group (median two days, interquartile range (IQR) one to three) compared to the M-TKA group (median four days, IQR three to five). The shorter length of stay observed in this study was considered a cost saving for the center relative to M-TKA.<sup>33</sup>

A prospective RCT carried out in Glasgow, UK, by Blythe et al.<sup>42</sup> compared M-PKA ( $n=65$ , mean age 62.5 years) to RA-PKA ( $n=64$ , mean age 62.1 years) at a single study center with patients enrolled between October 2010 and December 2012 and assessed through ten-year follow-up. Five patients in the M-PKA group were revised to TKA and there were no revisions at ten-years in the RA-PKA group, resulting in significantly greater re-intervention ( $p=0.001$ ), all-cause revision ( $p=0.007$ ) and aseptic revision ( $p=0.023$ ) for M-PKA compared to RA-PKA.<sup>42</sup> Median length of stay was two days for RA-PKA (IQR 2 to 3) and three days for M-PKA (IQR 2 to 4) ( $p=0.030$ ). The overall QALY gain per patient over the ten-year follow-up was greater in the RA-PKA group compared to M-PKA, though this did not achieve statistical significance (mean difference 0.186 (95% CI -0.626 to 0.999);  $p=0.651$ ).<sup>42</sup> The additional per patient costs associated with RA-PKA (£1,051) compared to M-PKA (£1850) resulted in a cost saving of £799 when accounting for all reinterventions. Two of the revisions in the M-PKA group were due to infection. When these two reinterventions were removed, RA-PKA was a cost-effective intervention with an incremental cost effectiveness ratio of £757 which was lower than the willingness-to-pay threshold (£20,000) suggested by the National Institute for Health and Care Excellence (NICE), and was cost-saving when more than 800 procedures per year are performed.<sup>42</sup>

Same day surgery for partial knee arthroplasty has been associated with potential cost benefits.<sup>67</sup> A study by Patel et al. evaluated Mako Partial Knee placed in an outpatient setting with same-day discharge from two outpatient Ambulatory Surgery Centers (ASCs).<sup>67</sup> A total of 63 patients underwent medial RA-PKA by one of six surgeons in the study.<sup>67</sup> Patients were surveyed preoperatively and at post-operative day one for their perception of outpatient joint replacement surgery preoperatively on a scale of 0 to 10 (0 negative to 10 positive).<sup>67</sup> Patient perception was positive throughout the process, with a preoperative mean perception of  $8.92 \pm 0.21$  and postoperative mean of  $9.3 \pm 0.2$ .<sup>67</sup> There were no complications or readmissions in the 30-day postoperative period.<sup>67</sup> However, one patient required a revision procedure within one year, and two patients required reoperation within one year, though none of these were attributed to aseptic loosening.<sup>67</sup> The revision was performed due to maltracking and the femoral component was removed and realigned and the polyethylene liner was exchanged.<sup>67</sup>

One reoperation was for a nerve ablation due to chronic pain in the operated knee, and the second reoperation was to remove excess cement from the joint space.<sup>67</sup> The authors conclude that outpatient robotic-arm assisted medial PKA was found to be feasible and was associated with a positive patient perspective of the procedure.

The cost-effectiveness studies described above all differed in inputs specific to their country, local region, hospital system or payer. These studies demonstrated that robotic-arm assisted partial knee arthroplasty, in comparison to manual TKA or manual partial knees, was associated with lower costs and/or improvements in QALY.<sup>33, 42, 46, 64-67</sup>

## 5. Conclusion

Mako Partial Knee offers the potential for surgeons to achieve component placement accuracy<sup>19</sup> and soft tissue balancing,<sup>20</sup> as well as to enhance clinical outcomes.<sup>12-14,32-38,48</sup> Patients have reported tangible benefits of robotic-arm assisted procedures, including treatment satisfaction,<sup>12,32,35,36</sup> return to activities of daily living<sup>51</sup> and a “forgotten” joint.<sup>13,32,33</sup> Surgeons are empowered to achieve their target preoperative plans with precision.<sup>19</sup> The cost-effectiveness studies described here demonstrated lower costs and better improvements in QALY for patients who received robotic-arm assisted partial knees in contrast to those received manual TKA or manual partial knees.<sup>65-66</sup> Ultimately, the potential benefits of Mako Partial Knee surgery are reported to be experienced by all key players – patients, surgeons and health systems.

## 6. References

1. Ollivier M, Abdel M, Parratte S, and Argenson JN. Lateral unicondylar knee arthroplasty (UKA): Contemporary indications, surgical technique, and results. *Int Orthop*. 2014;38(2):449–455.
2. Price AJ, Rees JL, Beard DJ, Gill RH, Dodd CA, Murray DM. Sagittal plane kinematics of a mobile bearing unicompartmental knee arthroplasty at 10 years: a comparative in vivo fluoroscopic analysis. *The Journal of Arthroplasty*. 2004;19(5):5907.
3. Schwab PE, Lavand'homme P, Yombi JC, Thienpont E. Lower blood loss after unicompartmental than total knee arthroplasty. *Knee surgery, sports traumatology, arthroscopy: official journal of the ESSKA*. 2015;23(12):3494-500.
4. Brown NM, Sheth NP, Davis K, Berend ME, Lombardi AV, Berend KR, et al. Total knee arthroplasty has higher postoperative morbidity than unicompartmental knee arthroplasty: a multicenter analysis. *The Journal of Arthroplasty*. 2012;27(8 Suppl):86-90.
5. Larsen K, Sorensen OG, Hansen TB, Thomsen PB, Soballe K. Accelerated perioperative care and rehabilitation intervention for hip and knee replacement is effective: a randomized clinical trial involving 87 patients with 3 months of follow-up. *Acta Orthopaedica*. 2008;79(2):149-59.
6. Jamali AA and Scott RD. Lateral unicompartmental knee arthroplasty. *Techniques in Knee Surgery*. 2005;4(2):79-88.
7. Pennington DW, Swienckowski JJ, Lutes WB, Drake GN. Lateral unicompartmental knee arthroplasty: survivorship and technical considerations at an average follow-up of 12.4 years. *J Arthroplasty*. 2006;21(1):13-17.
8. Volpi P, Marinoni L, Bait C, et al. Lateral unicompartmental knee arthroplasty: indications, technique and short-medium term results. *Knee Surg Sports Traumatol Arthrosc* 2007;15:1028–1034.
9. Odgaard A, Madsen F, Kristensen PW, Kappel A, and Fabrin J. The Mark Coventry Award: Patellofemoral Arthroplasty Results in Better Range of Movement and Early Patient-reported Outcomes Than TKA. *Clin Orthop Relat Res*. 2018; 476:87-100.
10. Citak M, Dersch K, Kamath AE, Haasper C, Gehrke T, Kendoff D. Common causes of failed unicompartmental knee arthroplasty: a single centre analysis of four hundred and seventy one cases. *Int Orthop*. 2014; 38:961-5.
11. Kazarian GS, Barrack TN, Okafor L, Barrack RL, Nunley RM, Lawrie CM. High Prevalence of Radiographic Outliers and Revisions with Unicompartmental Knee Arthroplasty. *The Journal of Bone and Joint Surgery. American Volume*. 2020 May 8;102:1151-9.
12. Kleeblad LJ, Borus TA, Coon TM, Douchis J, Nguyen JT, Pearle AD. Midterm Survivorship and Patient Satisfaction of Robotic-Arm-Assisted Medial Unicompartmental Knee Arthroplasty: A Multicenter Study. *The Journal of Arthroplasty*. 2018;33(6):1719-1726.
13. Blyth MJ, Anthony I, Rowe P, Banger MS, MacLean A, Jones B. Robotic arm-assisted versus conventional unicompartmental knee arthroplasty: Exploratory secondary analysis of a randomized controlled trial. *Bone and Joint Research*. 2017; 6(11):631-639.
14. Millar LJ, Banger M, Rowe PJ, Blyth M, Jones B, Maclean A. A five-year follow up of gait in robotic assisted vs conventional unicompartmental knee arthroplasty. *Gait & Posture*. 2018; 65 (Supplement 1): 31-32.
15. Australian Orthopaedic Association National Joint Replacement Registry (AOANJRR) Annual Report. 2024. [Available at <https://aoanjrr.sahmri.com/annual-reports-2024>]
16. NJR- The National Joint Registry. 2024. [Available from: <https://www.njr.org.uk/>]

17. Tsai TY, Dimitriou D, Liow MHL, Rubash HE, Li G, Kwon YM. Three-dimensional imaging analysis of unicompartmental knee arthroplasty evaluated in standing position: component alignment and in vivo articular contact. *J Arthroplasty*. 2016 May;31(5):1096-101.
18. Aleto TJ, Berend ME, Ritter MA, Faris PM, Meneghini RM. Early failure of unicompartmental knee arthroplasty leading to revision. *J Arthroplasty*. 2008 Feb;23(2):159-63.
19. Bell SW, Anthony I, Jones B, MacLean A, Rowe P, Blyth M. Improved accuracy of component positioning with robotic-assisted unicompartmental knee arthroplasty: data from a prospective, randomized controlled study. *J Bone and Joint Surg Am*. 2016;98(8):627-35.
20. Plate JF, Mofidi A, Mannava S, Smith BP, et al. Achieving accurate ligament balancing using robotic-assisted unicompartmental knee arthroplasty. *Advances in Orthopedics*. 2013;2013:837167.
21. St Mart J-P, Goh EL, Goudie E, Crawford R, English H, Donnelly W. Clinical and radiological outcomes of robotic-assisted unicompartmental knee arthroplasty: Early lessons from the first 100 consecutive knees in 85 patients. *The Knee*. 2022 Jan;34:195-205.
22. Kayani B, Konan S, Pietrzak JRT, Huq SS, Tahmassebi J, Haddad FS. The learning curve associated with robotic-arm assisted unicompartmental knee arthroplasty: a prospective cohort study. *Bone Joint J*. 2018 Aug;100-B(8):1033-1042.
23. Kazarian GS, Barrack RL, Barrack TN, Lawrie CM, Nunley RM. Radiographical outcomes following manual and robotic-assisted unicompartmental knee arthroplasty. *Bone Jt Open*. 2021;2(3):191-197.
24. Matassi F, Innocenti M, Giabbani N, Sani G, Cozzi Lepri A, Piolanti N, Civinini R. Robotic-assisted unicompartmental knee arthroplasty reduces components' positioning differences among high- and low-volume surgeons. *J Knee Surg*. 2022 Dec;35(14):1549-1555..
25. Gaggiotti S, Foissey C, Pineda T, Batailler C, Gaggiotti G, Gaggiotti S, Servien E, Lustig S. Enhancing robotic precision in medial UKA: Image-based robot-assisted system had higher accuracy in implant positioning than imageless robot-assisted system across 292 knees. *Knee Surg Sports Traumatol Arthrosc*. 2024 Aug;32(8):2097-2106.
26. Kumar A, Hung C-H, Hsieh S-L, Kuo C-C, Mao J-T, Lin E-T, et al. Makoplasty medial unicondylar knee replacement: Correction or postoperative angle matters? *Int J Med Robot*. 2022 Apr;18(2):e2356.
27. Jinnah R, Lippincott CJ, Horowitz S, Conditt MA. The learning curve of robotically assisted UKA. Paper No. 407, 56th Annual Meeting of the Orthopaedic Research Society. 69 March 2010
28. Tay ML, Carter M, Bolam SM, Zeng N, Young SW. Robotic-arm assisted unicompartmental knee arthroplasty system has a learning curve of 11 cases and increased operating time. *Knee Surgery, Sports Traumatology, Arthroscopy*. 2023 Mar;31(3):793-802.
29. Hampp E, Chang T-C, Pearle A. Robotic Partial Knee Arthroplasty Demonstrated Greater Bone Preservation Compared to Robotic Total Knee Arthroplasty. Orthopaedic Research Society 2019.
30. Hampp EL, Scholl L, Faizan A, Sodhi N, Mont MA, Westrich G. Comparison of iatrogenic soft tissue trauma in robotic-assisted versus manual partial knee arthroplasty. *Surg Technol Int*. 2021 Aug 5; 39:419-426..
31. Zuiderbaan HA, van der List JP, Khamaisy S, Nawabi DH, Thein R, Ishmael C, Paul S, Pearle AD. Unicompartmental knee arthroplasty versus total knee arthroplasty: Which type of artificial joint do patients forget? *Knee Surg Sports Traumatol Arthrosc*. 2017;25(3):681-686.
32. Clement ND, Bell A, Simpson P, Macpherson G, Patton JT, Hamilton DF. Robotic-assisted unicompartmental knee arthroplasty has a greater early functional outcome when compared to manual total knee arthroplasty for isolated medial compartment arthritis. *Bone & Joint Research*. 2020May;9(1):15-22.

33. van der List JP, Chawla H, Villa JC, Pearle AD. Different optimal alignment but equivalent functional outcomes in medial and lateral unicompartmental knee arthroplasty. *The Knee*. 2016;23(6):987-95.
34. Zuiderbaan HA, Khamaisy S, Thein R, Nawabi DH, Pearle AD. Congruence and joint space width alterations of the medial compartment following lateral unicompartmental knee. *Bone Joint J*. 2015. 97-B(1): 50-5.
35. Thein R, Zuiderbaan HA, Khamaisy S, Nawabi DH, Poultsides LA, Pearle AD. Medial Unicompartmental Knee Arthroplasty Improves Patellofemoral Congruence: A Possible Mechanistic Explanation for Poor Association Between Patellofemoral Degeneration and Clinical Outcome. *J Arthroplasty*. 2015;30(11):1917-22.
36. Khamaisy S, Zuiderbaan HA, van der List JP, Nam D, Pearle AD. Medial unicompartmental knee arthroplasty improves congruence and restores joint space width of the lateral compartment. *The Knee*. 2016; 23(3):501-505.
37. Roche M, Vakharia RM, Law T, Rodriguez HC, Mekkawy KL, Sodhi N, Dubin JA, Hameed D, Mont MA. Excellent 10-Year survivorship of robotic-arm-assisted unicompartmental knee arthroplasty. *Journal of Orthopaedics*. 2024 Mar;55:32-37.
38. Burger JA, Kleeblad LJ, Laas N, Pearle AD. Mid-term survivorship and patient-reported outcomes of robotic-arm assisted partial knee arthroplasty: a single-surgeon study of 1,018 knees. *The Bone & Joint Journal*. 2020 Jan;102-B(1):108-16.
39. Pearle AD, van der List JP, Lee L, Coon TM, Borus TA, Roche MW. Survivorship and patient satisfaction of robotic-assisted medial unicompartmental knee arthroplasty at a minimum two-year follow-up. *Knee*. 2017;24(2):419-428.
40. Gilmour A, MacLean AD, Rowe PJ, Banger MS, Donnelly I, Jones BG, Blyth MJG. Robotic-Arm-Assisted vs Conventional Unicompartmental Knee Arthroplasty. The 2-Year Clinical Outcomes of a Randomized Controlled Trial. *The Journal of Arthroplasty*. 2018;33(7S):S109-S115.
41. Banger M, Doonan J, Rowe P, Jones B, MacLean A, Blyth MJB. Robotic arm-assisted versus conventional medial unicompartmental knee arthroplasty: five-year clinical outcomes of a randomized controlled trial. *Bone Joint J* 2021;103-B(6):1088-1095
42. Blyth MJG, Clement ND, Choo XY, Doonan J, MacLean A, Jones BG. Robotic arm-assisted medial compartment knee arthroplasty is a cost-effective intervention at ten-year follow-up. *Bone Joint J*. 2025;107-B(1):72-80.
43. Bayoumi T, Kleeblad LJ, Borus TA, Coon TM, Douchis J, Nguyen JT, Pearle AD. Ten-Year Survivorship and Patient Satisfaction Following Robotic-Arm-Assisted Medial Unicompartmental Knee Arthroplasty: A Prospective Multicenter Study. *J Bone Joint Surg Am*. 2023 Jun 21;105(12):933-942.
44. Zambianchi F, Seracchioli S, Franceschi G, Cuoghi Costantini R, Malatesta A, Barbo G, Catani F. Image-based robotic-arm assisted unicompartmental knee arthroplasty provides high survival and good-to-excellent clinical outcomes at minimum 10 years follow-up. *Knee Surg Sports Traumatol Arthrosc*. 2023 Dec;31(12):5477-5484.
45. Heckmann ND, Antonios JK, Chen XT, Kang HP, Chung BC, Piple AS, et al. Midterm Survivorship of Robotic-Assisted Lateral Unicompartmental Knee Arthroplasty. *The Journal of Arthroplasty*. 2022 May;37(5):831-6.
46. Cool CL, Needham KA, Khlopas A, Mont MA. Revision Analysis of Robotic Arm-Assisted and Manual Unicompartmental Knee Arthroplasty. *J Arthroplasty*. 2019;34(5):926-931. doi:10.1016/j.arth.2019.01.018
47. St Mart J-P, de Steiger RN, Cuthbert A, Donnelly W. The three-year survivorship of robotically assisted versus non-robotically assisted unicompartmental knee arthroplasty: a study from the Australian Orthopaedic Association National Joint Replacement Registry. *The Bone & Joint Journal*. 2020 Mar;102-B(3):319-28.
48. Dretakis K, Igoumenou VG. Outcomes of robotic-arm-assisted medial unicompartmental knee arthroplasty: minimum 3-year follow-up. *Eur J Orthop Surg Traumatol*. 2019;29(6):1305-1311.
49. Deese JM, Gratto-Cox G, Carter DA, et al. Patient reported and clinical outcomes of robotic-arm assisted unicompartmental knee arthroplasty: minimum two year follow-up. *J Orthop*. 2018 Aug 16;15(3):847-853.

50. Swedish knee registry 2024 [report available from <https://sar.registercentrum.se/>]
51. Coon T, Shi S, DeBattista J. Clinical and functional outcomes of robotic arm assisted medial unicompartmental knee arthroplasty. European Knee Society 2017 Annual Meeting. London, England. Poster No. P59. April 19-21, 2017.
52. Zambianchi F, Daffara V, Franceschi G, Banchelli F, Marcovigi A, Catani F. Robotic arm-assisted unicompartmental knee arthroplasty: high survivorship and good patient-related outcomes at a minimum five years of follow-up. *Knee Surg Sports Traumatol Arthrosc.* 2021;29(10):3316-3322. doi:10.1007/s00167-020-06198-9
53. Motesharei A, Rowe P, Blyth M, Jones B, Maclean A. A comparison of gait one year post operation in an RCT of robotic UKA versus traditional Oxford UKA. *Gait & Posture.* 2018; 62:41–45.
54. Borus T, Roberts D, Fairchild P, Pirtle K, Baer M. Early Functional Performance of Unicompartmental Knee Arthroplasty Compared to Total Knee Arthroplasty. 2nd World Arthroplasty Congress (WAC) 2018. Rome, Italy. ePoster P4. April 19-21, 2018.
55. Coon T, Shi S, DeBattista J, Bhowmik-Stoker M. Clinical and functional outcomes of robotic arm assisted unicompartmental and bicompartamental knee arthroplasty. European Knee Society 2017 Annual Meeting. London, England. Poster No. P60. April 19-21, 2017.
56. Augart MA, Plate JF, Bracey DN, Jinnah A, Poehling GG, Jinnah RH. Robotic Lateral and Medial Unicompartmental Knee Arthroplasty. *Operative Techniques in Orthopaedics.* 2015;25(2):95-103.
57. Burger JA, Dooley MS, Kleeblad LJ, Zuiderbaan HA, Pearle AD. What is the impact of patellofemoral joint degeneration and malalignment on patient- reported outcomes after lateral unicompartmental knee arthroplasty? *The Bone & Joint Journal.* 2020 Jun;102-B(6):727-35.
58. Farr J and Barrett D. Optimizing patellofemoral arthroplasty. *Knee.* 2008;15(5):339-47.
59. Noyes FR, Fleckenstein CM, Nolan J. Return to Sports in Patients Aged 50 Years or Younger After Robotic-Assisted Patellofemoral Arthroplasty: A 10-Year Experience Reporting High Clinical Benefits and High Patient Satisfaction With Return to an Active Lifestyle. *The American Journal of Sports Medicine.* 2024;52(6):1514-1526.
60. Lonner JH. Modular bicompartamental knee arthroplasty with robotic arm assistance. *Am J Orthop (Belle Mead NJ).* 2009 Feb;38(2 Suppl):28-31.
61. Thienpont E, Price A. Bicompartamental knee arthroplasty of the patellofemoral and medial compartments. *Knee Surg Sports Traumatol Arthrosc.* 2013 Nov;21(11):2523-31.
62. Gaudiani MA, Samuel LT, Diana JN, DeBattista JL, Coon TM, Moore RE, Kamath AF. Robotic-arm assisted bicompartamental knee arthroplasty: durable results up to 7-year follow-up. *Int J Med Robot.* 2022 Feb;18(1):e2338.
63. Sun Y, Liu W, Hou J, Hu X, Zhang W. Does robotic-assisted unicompartmental knee arthroplasty have lower complication and revision rates than the conventional procedure? A systematic review and meta- analysis. *BMJ Open.* 2021 Aug;11(8):e044778.
64. Kazarian GS, Lonner JH, Maltenfort MG, Ghomrawi HMK, Chen AF. Cost-Effectiveness of Surgical and Nonsurgical Treatments for Unicompartmental Knee Arthritis: A Markov Model. *J Bone Joint Surg Am* 2018;100:1653-60.
65. Clement ND, Deehan DJ, Patton JT. Robot-assisted unicompartmental knee arthroplasty for patients with isolated medial compartment osteoarthritis is cost-effective: A MARKOV DECISION ANALYSIS. *Bone Joint Journal* 2019;101-B(9):1063–1070.
66. Varughese I, Whitehouse SL, Donnelly WJ, Crawford RW. The Cost Effectiveness of Unicompartmental versus Total Knee Arthroplasty. *J Knee Surg.* 2024 Jan;37(1):66-72.
67. Patel T, Baker J, Eberhard A, Ginesin E, Patel RM. Robotic-Assisted Medial Unicompartmental Knee Arthroplasty in an Outpatient Setting. *Surg Technol Int.* 2025 Jan 29;45:sti45/1844. Epub ahead of print.

---

A surgeon must always rely on his or her own professional clinical judgment when deciding whether to use a particular product when treating a particular patient. Stryker does not dispense medical advice and recommends that surgeons be trained in the use of any particular product before using it in surgery.

The information presented is intended to demonstrate the breadth of Stryker's product offerings. A surgeon must always refer to the package insert, product label and/or instructions for use before using any of Stryker's products. Products may not be available in all markets because product availability is subject to the regulatory and/or medical practices in individual markets. Please contact your sales representative if you have questions about the availability of products in your area.

Stryker Corporation or its divisions or other corporate affiliated entities own, use or have applied for the following trademarks or service marks: Mako, Stryker. All other trademarks are trademarks of their respective owners or holders.

**JR-MPKA-OTHD-647153\_REV-1**  
Copyright © 2026 Stryker

**Stryker**  
325 Corporate Drive  
Mahwah, NJ 07430  
t: 201 831 5000  
stryker.com